

| | Raptor Species | Sun | Mon | Tue | Wed | Thu | Fri | Sat |
|----------------------------|-----------------------|-----------------|-----|----------------|-----|---------------------------|-----|-------------------------|
| Feed during daylight hours | Red-tailed Hawk | 1 Q | 4 M | 4 C | 1 R | 1 Q | 4 C | 1 R |
| | American Goshawk | 1 Q | 4 M | 4 C | 1 R | 1 Q | 4 C | 1 R |
| | Cooper's Hawk | 2 C | 3 M | 3 C | ½ Q | ½ Q | 2 C | 3 M |
| | Sharp-shinned Hawk | 2 C | 2 M | 2 C | 2 M | 2 C | 2 M | 2 C |
| | Broad-winged Hawk | 2 C | 3 M | 3 C | ½ Q | ½ Q | 2 C | 1 R |
| | Red Shouldered Hawk | 3 C | 3 M | 3 C | ½ Q | ½ Q | C | 1 R |
| | Rough-legged Hawk | 1 Q | 4 M | 4 C | 1 R | 1 Q | 4 C | 1 R |
| | Swainson's Hawk | 1 Q | 4 M | 4 C | 1 R | 1 Q | 4 C | 1 R |
| | American Kestrel | 1 C | 2 M | 2 C | 1 M | 1 C | 2 C | 2 M |
| | Merlin | 2 C | 2 M | ½ Q | ½ Q | 2 M | 2 C | 2 M |
| | Peregrine Falcon | 4 C | 1 Q | 4 C | 1 Q | 5 C | 1 Q | 4 C |
| | Gyr Falcon | 5 C | 1 Q | 1 R | 1 Q | 5 C | 1 Q | 1 Q |
| | Prairie Falcon | 4 C | ½ Q | ½ Q | 4 M | 4 C | ½ Q | ½ Q |
| | Turkey Vulture | 1 R | 6 C | 1 R | 1 Q | 5 C | 6 M | 1 Q |
| | Bald Eagle | 2 Q | 1 R | 2 Q | 1 F | 2 R | 2 Q | 1 F |
| Golden Eagle | 2 Q | 1 R | 2 Q | 1 R | 2 R | 2 Q | 1 R | |
| Feed after dusk | Great Horned Owl | 1 Q | 4 M | 4 C | 1 R | 1 Q | 4 C | 1 R |
| | Barred Owl | 1 Q | 3 M | 4 C | 1 R | 1 Q | 3 C | 1 R |
| | Barn Owl | 1 Q | 4 M | 1 R | 4 M | 1 R | 4 C | 1 R |
| | Eastern Screech-owl | 1 C | 2 M | 2 C | 1 M | 1 C | 2 C | 2 M |
| | Northern Saw-whet Owl | 1 C | 2 M | 2 C | 1 M | 1 C | 2 C | 2 M |
| Q = Quail | | M = Mice | | R = Rat | | C = Day Old Chicks | | F = Fish (11 OZ) |

Notes:

If necessary, temporarily substitute: 1 Q with 1 R, 1 M with 1 C, 1 F with 2 Q
Purchase medium rats, large mice, and 8-week-old quail.